
Unclassified
SECURITY CLASSIFICATION OF THIS PAGE

DTIC FILE COPY



REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188
1a REPORT SECURITY CLASSIFICATION Unclassified	16. RESTRICTIVE MARKINGS				
2a. SECURITY CLASSIFICATION A THE STY AUG 22 1990 2b. DECLASSIFICATION / DOWNGRAD 5 SCHEDULE		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release and sale; distribution unlimited.			
4. PERFORMING ORGANIZATION REPORT NUM E (S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)			
Report #7		#4135010			
6a. NAME OF PERFORMING ORGANIZATION 6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION			
University of Minnesota		Office of Naval Research			
6C ADDRESS (City, State, and ZIP Code)		7b. ADDRESS (City, State, and ZIP Code)			
Department of Chemistry					
207 Pleasant Street SE		300 N. Quincy Street			
Minneapolis, MN 55455-0431		Arlington, VA 22217-5000			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			
Office of Naval Research	N0001489J1301				
8c. ADDRESS (City, State, and ZIP Code)		10 SOURCE OF FUNDING NUMBERS			
800 N. Quincy Street		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
Arlington, VA 22217-5000		61153N	4135	110.	4135010
11. TITLE (Include Security Classification)		0113311	4133	l	1425510
,					
"Symmetry Bias in H-Bonded Crystals"					
12. PERSONAL AUTHOR(S) 11.C. Etter, and K.S. Huang					
13a. TYPE OF REPORT 13b. TIME CO	VEDCO				
Technical FROM	May 23, 1	4 DATE OF REPORT (Year, Month, Day) 15 PAGE COUNT 11ay 23, 1990 2			
16 SUPPLEMENTARY NOTATION					
To be published in Chemistry of 'Laterials					
17. COSATI CODES 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)					
FIELD GROUP SUB-GROUP					
37 03 576 2					
19 ABSTRACT (Continue on reverse if necessary and identity by block number)					
The Cambridge Crystallographic Data Base (CCDB) was searched for all examples of small molecules					
with nitroaniline groups. This data set was divided into primary aned secondary nitroanilines (32 compounds),					
tertiary nitroanilines (36 compounds). For each set we determined what percent of the crystal structures were					
acentric. For the primary and secondary nitroanilines, which can form nitroaniline hydrogen bonds, 41% of the					
crystals were acentric. For the tertiary nitroanilines, without -NH2O ₂ N hydrogen bonds, 17% were acentric.					
These numbers are to be compared to the frequency of occurrence of acentricity in the CCDB for the total					
population of all organic molecules. This number was determined several years ago when the data base containe about 40,000 structures. At that time 25% of the structures were acentric. Within that set it is estimated that most					
of the acentric structures contain resolved chiral compounds. Thus, a better, yet generous, estimate for our					
purposes would be that only about 10% of all racemic or achiral compounds pack in acentric crystal structures.					
Thus, there appears to be a significant bias towards crystallographic acentricity for the primary and secondary					
nitroaniline structures. Further work is in progress to determine if the hydrogen-bond patterns in the primary and					
secondary nitroaniline class are themselves acentric.					
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT 21 ABSTRACT SECURITY CLASSIFICATION					
■ UNCLASSIFIED/UNLIMITED □ SAME AS RP	Unclassified				
224 NAME OF RESPONSIBLE INDIVIDUAL	T DTIC USERS	226 TELEPHONE (IF			
DD Form 1473, JUN 86	Previous editions are o	hsolete	CECLIBITY (LASSIEICA	TION OF THIS PAGE

Examples of Acentric Primary Nitroanilines and Centric Tertiary Analogs



may be biasing the final crystal structures to be acentric. Dipole-dipole interactions Both acentric compounds pack with acentric hydrogen-bond networks which will be a less important contributor to the final packing pattern in structures with hydrogen-bonded networks than in structures without, so acentricity may be seen more frequently.

Availability Codes

Distribution/

Arail and/or

Accession For

GRALI

Justiftention.

Ur.aru.ormoed

DIIC TAB

Our results show that for primary and secondary nitroanilines, acentric crystal structures occur about 40% of the time, while for tertiary nitroanilines acentricity occurs 17% of the time (about the same rate as for other achiral molecules),